




This document is provided by NHTSA in an effort to help State EMS Officials track particular EMS activations occurring during the COVID-19 pandemic. This document will be updated periodically to provide current information regarding temporal variations in the type and characteristics of EMS activations occurring in the U.S. during the COVID-19 outbreak.



When assessing data contained in the National EMS Repository, it is important to recognize that there can be a lag in the submission of patient care reports to the Repository. Looking in the lower left-hand corner, the definition of “Submission Lag” is the date/time difference (in hours) between the completion of an EMS activation (eTimes.13 - Unit Back in Service Date/Time) and the arrival date/time of the record in the National EMS Repository.

- 25 hours ~ 1 day
- 60 hours ~ 2.5 days
- 177 hours ~ 7 days

Thus, the generalizability of a “count” or “rate” associated with any week or day should be assessed in light of the completeness of data for that date.

Identification of ILI

- EMS Primary and Secondary Impression
 - B79 codes: **SARS and other coronavirus**
 - J09 codes: **Influenza**
 - J15 codes: **Pneumonia**
- Patient Primary and Associated Symptoms
 - R05 codes: **Cough**
 - R06 codes: **Shortness of Breath**
 - R50 codes: **Fever**
 - J02 codes: **Pharyngitis**

The definition of Influenza-Like Illness (ILI) is based on the record inclusion criteria provided in the User Guide for the National NEMESIS ILI Surveillance Dashboard. The ILI Surveillance Dashboard User Guide can be found at: <https://wiki.utahdcc.org/confluence/x/BAKXAg>.

Provided in this slide are examples of the two types of ICD-10-CM codes included in the ILI criteria.




Research Sample

- Time Period 1 (40th week of 2017 to the 26th week of 2018): **10,205,536**
- Time Period 2: (40th week of 2018 to the 26th week of 2019): **15,588,588**
- Time Period 3: (40th week of 2019 to the 26th week of 2020): **17,811,401**




Three date/time samples of EMS activations are included in this assessment of the COVID-19 pandemic. Two date/time periods (from the 40th week [late September] of the previous year to the 26th week [last full week of June] of the next year) are included to provide reference comparison to the third time period of interest (the 40th week of 2019 through the 26th week of 2020). The total sample includes 43,605,525 9-1-1 initiated EMS activations resulting in patient contact.





The number of States submitting to the National EMS Repository increased over the study period (2017 – 32 States, 2018 – 40 States, 2019 – 44 States). The District of Columbia submitted PCR's in each time period. States enrolling in the National EMS Repository commonly begin submitting PCR's at the beginning of the calendar year. No state has stopped submitting PCR's once enrolled.

The number of EMS activations decreased by approximately 34% between Week 10 (March 2nd to March 8th) and Week 17 (April 20th through April 26th 2020). The number of EMS activations began a second downward trend in Week 20.




The count of EMS activations related to ILI symptoms, greatly increased beginning in Week 10 through Week 14, with a dramatic precipitous drop thereafter through Week 24. Because of the variation in the count of EMS activations through time, rates were calculated to remove difference due solely the count of submitted records.







The count of cardiac arrest is created by summing the NEMESIS element eArrest.01 - Cardiac Arrest: Yes, Prior to EMS Arrival AND Yes, After EMS Arrival. The increased number of EMS attended cardiac arrests from week 10 through week 15 represent approximately 1,000 additional cardiac arrests.




The count of EMS attended scene deaths results from summing the NEMSIS element eDisposition.12 - Incident/Patient Disposition: Patient Dead at Scene - With (or Without) Resuscitation Attempted and With (or Without) Transport.




Similar date stamps are superimposed across the dramatic shifts in rate of EMS attended scene deaths.




The rate of possible injury-related EMS activations demonstrates the expected increase during warmer months, but lower than expected rates beginning in Week 10 and beginning to increase in Week 13. Similar date stamps are superimposed across the dramatic shifts in rate of EMS activations reporting a possible injury.



The count of EMS activations related to vehicle crashes results from summing the NEMSIS element eDispatch.01 - Complaint Reported by Dispatch: Traffic/Transportation Incident.



Similar date stamps are superimposed across the dramatic shifts in rate of EMS activations associated with a Traffic/Transportation Incident.




The count of opioid-related EMS activations results from summing the NEMSIS elements: eSituation.11 - Provider's Primary Impression, AND eSituation.12 - Provider's Secondary Impressions, AND eSituation.09 - Primary Symptom, AND eSituation.10 - Other Associated Symptoms with any of the following ICD-10-CM codes:


Codes

F11 codes – Opioid-related disorders


T40 codes - Poisoning by (and adverse effects of) opioid-related drugs




Similar date stamps are superimposed across shifts in the rate of EMS activations documenting opioid-related issues.



The count of EMS activations with documented use of Naloxone results from summing the NEMSIS element: eProcedures.03 – Procedures performed on the Patient. An EMS activation indicating multiple administrations of Naloxone to the same patient is counted only once.



Similar date stamps are superimposed across shifts in the rate of EMS activations with documented Naloxone use.



The count of EMS activations related to suicide/self-harm results from summing the NEMSIS elements: eSituation.11 - Provider's Primary Impression, AND eSituation.12 - Provider's Secondary Impressions, AND eSituation.09 - Primary Symptom, AND eSituation.10 - Other Associated Symptoms with any of the following ICD-10-CM codes:

R45 codes - Suicidal ideations


T14.91 - Suicide attempt

T40 codes - Poisoning by medicaments, intentional self-harm


T50 codes - Poisoning by unspecified drugs, medicaments and biological substances, intentional self-harm

T65 codes - Toxic effect of specified and unspecified substances, intentional self-harm

X71 through X83 codes - Intentional self-harm by specified and unspecified means




Similar date stamps are superimposed across shifts in the rate of EMS activations documenting suicide/self harm issues.




The count of EMS activations documenting alcohol-related issues as an impression or symptom results from summing the NEMSIS elements: eSituation.11 - Provider's Primary Impression, AND eSituation.12 - Provider's Secondary Impressions, AND eSituation.09 - Primary Symptom, AND eSituation.10 - Other Associated Symptoms with any of the following ICD-10-CM codes:

F10 codes – Alcohol-related disorders




Similar date stamps are superimposed across shifts in the rate of EMS activations documenting alcohol-related issues as an impression or symptom.



The count of EMS activations related to mental/behavioral health issues results from summing the NEMSIS elements: eSituation.11 - Provider's Primary Impression, AND eSituation.12 - Provider's Secondary Impressions, AND eSituation.09 - Primary Symptom, AND eSituation.10 - Other Associated Symptoms with any of the following ICD-10-CM codes:

- F41.9 – Anxiety, NOS
- F41.1 - Generalized anxiety disorder
- R41.82 - Altered mental status, unspecified
- F32.9 - Major depression, NOS
- F99 - Mental disorder, NOS
- R45.89 - Other symptoms and signs involving emotional state
- R45.7 - State of emotional shock and stress, unspecified
- R46.2 - Strange and inexplicable behavior
- R46 - Symptoms and signs involving appearance and behavior
- R45.82 - Worries



Similar date stamps are superimposed across shifts in the rate of EMS activations associated with mental/behavioral health issues.

The image shows a screenshot of the NEMESIS website homepage. At the top left is the NEMESIS logo with the tagline "BETTER DATA. BETTER CARE." Below the logo is a navigation menu with links for "WHAT IS NEMESIS", "USING EMS DATA", "VIEW REPORTS", "CALLS AND TRAININGS", "TECHNICAL RESOURCES", and "SUPPORT". A search bar is located below the navigation menu. The main content area features a large banner with a hand holding a glowing 3D data cube. To the right of the cube, the text reads "NEMESIS V3 EMS Data Cube Now available!". In the bottom left corner of the screenshot is the "ems.gov" logo. Overlaid on the top half of the screenshot is the text "Questions?" in a large blue font, followed by "www.nemesis.org" in a smaller black font.

Please contact the NEMESIS Technical Assistance Center for updates to this document.
Contact N. Clay Mann at clay.mann@hsc.utah.edu.